DECLINE AND FALL OF THE STANDARD MODEL?

JOHN ELLIS
Theoretical Physics Division, CERN, CH-1211 Geneva 23
E-mail: John.Ellis@cern.ch

arXiv:hep-ph/0110192v1 15 Oct 2001

Motivations for physics beyond the Standard Model are reviewed, with particular emphasis on supersymmetry at the TeV scale. Constraints on the minimal supersymmetric extension of the Standard Model with universal soft supersymmetry-breaking terms (CMSSM) are discussed. These are also combined with the supersymmetric interpretation of the anomalous magnetic moment of the muon. The prospects for observing supersymmetry at accelerators are reviewed using benchmark scenarios to focus the discussion. Prospects for other experiments are discussed, including the detection of cold dark matter, $\mu \to e\gamma$ and related processes, as well as proton decay.

CERN-TH/2001-275 hep-ph/0110192

1 Introduction

The empire of the Standard Model has resisted all attacks by accelerator data. Nevertheless, we theorists are driven to overcome our ignorance of the barbarian territory beyond its frontiers. In the gauge sector, the Standard Model has three independent gauge couplings and (potentially) a CP-violating phase in QCD. In the Yukawa sector, it has six random-seeming quark masses, three charged-lepton masses, three weak mixing angles and the Kobayashi-Maskawa phase. Finally, the symmetry-breaking sector has at least two free parameters. Moreover, this list of 19 parameters in the Standard Model begs the more fundamental questions of the origins of the particle quantum numbers. As if this were not enough, non-accelerator neutrino experiments [1] now convince us that we need three neutrino mass parameters, three neutrino mixing angles and three CP-violating phases in the neutrino sector: one observable in oscillation experiments and two that affect $\beta\beta_{0\nu}$ experiments, without even talking about the mechanism of neutrino mass generation. Moreover, we should not forget about gravity, with at least two parameters to understand: Newton's constant $G_N \equiv m_P^{-2} \sim (10^{19}\ \mathrm{GeV})^{-2}$ and the cosmological 'constant', which recent data suggest is nonzero [2], and may not even be constant. Talking of cosmology, we would need at least one extra parameter to produce an inflationary potential, and at least one other to generate the baryon asymmetry, which cannot be explained within the Standard Model.

Confronted by our ignorance of so much barbarian territory, we legions of theorists organize our explorations on three main fronts: *unification* – the quest for a single framework for all gauge interactions, *flavour* – the quest for explanations of the proliferation of quark and lepton types, their mixings and CP violating phases, and *mass* – the quest for the origin of particle masses and an explanation why they are so much smaller than the Planck mass $m_P \sim 10^{19}$ GeV. Beyond all these beyonds, other scouting parties of theorists seek a *Theory of Everything* that includes gravity, reconciles it with quantum mechanics, explains the origin of space-time and why we live in four dimensions (if we do so).

Physics beyond the Standard Model is therefore a very broad subject. However, many aspects are discussed here by other speakers: electroweak flavour physics [3], CP violation [4], the Higgs sector [5], $g_\mu - 2$ [6], searches for new particles [7], neutrinos [8], dark

matter [9], strings and extra dimensions [10]. Therefore, in this talk I seek a complementary approach.

For reasons that I describe in Section 2, many theorists believe that supersymmetry is the inescapable framework for discussing physics at the TeV scale and beyond. In the rest of this talk, I first discuss the constraints imposed on (the simplest) supersymmetric models by the available experimental and cosmological constraints, then address the prospects for understanding $g_\mu - 2$ in supersymmetric models, the prospects for detecting sparticles directly at present and future colliders, and the prospects for non-collider experiments, including the searches for dark matter, $\mu \to e\gamma$ and proton decay.



Figure 1. Predictions for the radiative corrections ϵ_i in the Standard Model and a minimal one-generation model [14] are compared with the precision electroweak data [13].

2 The Electroweak Vacuum

The generation of particle masses requires the breaking of gauge symmetry in the vacuum:

$$m_{W,Z} \neq 0 \Leftrightarrow < 0|X_{I,I_3}|0 > \neq 0 \qquad (1)$$

for some field X with isospin I and third component I_3. The measured ratio

$$\rho \equiv \frac{m_W^2}{m_Z^2 \, \cos^2\theta_W} \simeq 1 \qquad (2)$$

tells us that X mainly has $I = 1/2$ [11], which is also what is needed to generate fermion masses. The key question is the nature of the field X: is it elementary or composite? A fermion-antifermion condensate $v \equiv < 0|X|0 > = < 0|\bar{F}F|0 > \neq 0$ would be analogous to what we know from QCD, where $< 0|\bar{q}q|0 > \neq 0$, and conventional superconductivity, where $< 0|e^-e^-|0 > \neq 0$. However, analogous 'technicolour' models of electroweak symmetry breaking [12] fail to fit the values of the radiative corrections ϵ_i to ρ and other quantities extracted from the precision electroweak data provided by LEP and other experiments, as seen in Fig. 1 [13]. One cannot exclude the possibility that some calculable variant of technicolour might emerge that is consistent with the data, but for now we focus on elementary Higgs models.

Within this framework, the data favour a relatively light Higgs boson, with $m_H \simeq 115$ GeV, just above the exclusion unit provided by direct searches at LEP, being the 'most-probable' [15]. This is one reason why many theorists were excited by the possible sighting during the last days of LEP of a Higgs boson, with a preferred mass of 115.6 GeV [7]. If this were to be confirmed, it would suggest that the Standard Model breaks down at some relatively low energy $\lesssim 10^3$ TeV [16]. As seen in Fig. 2, above this scale the effective Higgs potential of the Standard Model becomes unstable as the quartic Higgs self-coupling is driven negative by radiative corrections due to the relatively heavy top quark [17]. This is not necessarily a disaster, and it is possible that the present electroweak vacuum might be metastable, provided that its lifetime is longer than the age of the Universe [18]. However, we would surely feel more secure if such instability could be avoided.



Figure 2. The range allowed for the mass of the Higgs boson if the Standard Model is to remain valid up to a given scale Λ. In the upper part of the plane, the effective potential blows up, whereas in the lower part the present electroweak vacuum is unstable [17].



Figure 3. (a) If the quartic coupling M_0 (3) is too large, the effective potential blows up (solid line), whereas it is unstable if M_0 is too small (dotted line), indicating a need for fine tuning. (b) This occurs naturally in a supersymmetric model (solid line) but not if the $\tilde{H}$ are omitted (dotted line) [16].

This may be done by introducing new bosons ϕ coupled to the Higgs field [16]:

$$\lambda_{22}|H|^2\ |\phi|^2 \quad : \quad M_0^2 \equiv \lambda_{22} v^2 \qquad (3)$$

As seen in Fig. 3a, the effective potential is very sensitive to the coupling parameter M_0: for $M_0 \leq 70.9$ GeV in this example, the potential still collapses, whereas for $M_0 \geq 71.0$ GeV the potential blows up instead. Thus the bosonic coupling (3) must be finely tuned [16]. This occurs naturally in supersymmetry, in which the Higgs bosons are accompanied by fermionic partners $\tilde{H}$. As seen in Fig. 3b, again the Higgs coupling blows up in the absence of the $\tilde{H}$, whereas it is well behaved in the minimal supersymmetric extension of the Standard Model (MSSM).

The avoidance of fine tuning has long been the primary motivation for supersymmetry at the TeV scale [19]. This issue is normally formulated in connection with the hierarchy problem: why/how is $m_W \ll m_P$, or equivalently why is $G_F \sim 1/m_W^2 \gg G_N = 1/m_P^2$, or equivalently why does the Coulomb potential in an atom dominate over the Newton potential, $e^2 \gg G_N m_p m_e \sim (m/m_P)^2$, where $m_{p,e}$ are the proton and electron masses? One might think naively that it would be sufficient to set $m_W \ll m_P$ by hand. However, radiative corrections tend to destroy this hierarchy. For example, one-loop diagrams generate

$$\delta m_W^2 = \mathcal{O}\left(\frac{\alpha}{\pi}\right)\ \Lambda^2 \gg m_W^2 \qquad (4)$$

where Λ is a cut-off representing the appearance of new physics, and the inequality in (4) applies if $\Lambda \sim 10^3$ TeV, and even more so if $\Lambda \sim m_{GUT} \sim 10^{16}$ GeV or $\sim m_P \sim 10^{19}$ GeV. If the radiative corrections to a physical quantity are much larger than its measured values, obtaining the latter requires strong cancellations, which in general require fine tuning of the bare input parameters. However, the necessary cancellations are natural in supersymmetry, where one has equal numbers of bosons B and fermions F with equal couplings, so that (4) is replaced by

$$\delta m_W^2 = \mathcal{O}\left(\frac{\alpha}{\pi}\right)\ |m_B^2 - m_F^2|\ . \qquad (5)$$

The residual radiative correction is naturally small if

$$|m_B^2 - m_F^2| \lesssim 1 \text{ TeV}^2 \qquad (6)$$

Note that this argument is logically distinct from that in the previous paragraph. There supersymmetry was motivated by the control of logarithmic divergences, and here by the absence of quadratic divergences.

3 The MSSM

The MSSM has the same gauge interactions as the Standard Model, and similar Yukawa couplings. A key difference is the necessity of two Higgs doublets, in order to give masses to all the quarks and leptons, and to cancel triangle anomalies. This duplication is important for phenomenology: it means that there are five physical Higgs bosons, two charged $H^\pm$ and three neutral h, H, A. Their quartic self-interactions are determined by the gauge interactions, solving the vacuum instability problem mentioned above and limiting the possible mass of the lightest neutral Higgs boson. However, the doubling of the Higgs multiplets introduces two new parameters: $\tan\beta$, the ratio of Higgs vacuum expectation values and μ, a parameter mixing the two Higgs doublets.

There are two key experimental hints in favour of supersymmetry. One is provided by the LEP measurements of the gauge couplings, that are in very good agreement with supersymmetric GUTs [20] if sparticles weigh ~ 1 TeV. This agreement appears completely fortuitous in composite Higgs models [12], and is difficult (though not impossible [21]) to reproduce accurately in models with large extra dimensions [22]. The other experimental hint is provided by the preference of the precision electroweak data for a relatively light Higgs boson [15]. In the MSSM, one predicts $m_h \lesssim 130$ GeV [23,5], right in the preferred range, whereas composite Higgs model generally predict heavier effective Higgs masses.

The gauge symmetries of the MSSM would permit the inclusion of interactions that violate baryon number and/or lepton number [24]:

$$\lambda LLE^c + \lambda' QD^c L + \lambda'' U^c D^c D^c \qquad (7)$$

where the $L(Q)$ are left-handed lepton (quark) doublets and the $E^c(D^c, U^d)$ are conjugates of the right-handed lepton (quark) singlets. Their possible appearance is ignored in this talk, in which case the lightest supersymmetric particle is stable, and hence a candidate for dark matter [25]. In the following this is assumed to be a neutralino, i.e., a mixture of the $\tilde{\gamma}, \tilde{H}$ and $\tilde{Z}$.

The final ingredient in the MSSM is the soft supersymmetry breaking, in the form of scalar masses m_0, gaugino masses $m_{1/2}$ and trilinear couplings A [26]. These are presumed to be inputs from physics at some high-energy scale, e.g., from some supergravity or superstring theory, which then evolve down to lower energy scale according to well-known renormalization-group equations. In the case of the Higgs multiplets, this renormalization can drive the effective mass-squared negative, triggering electroweak symmetry weaking [27]. In this talk, it is assumed that the m_0 are universal at the input scale [a], as are the $m_{1/2}$ and A parameters. In this case the free parameters are

$$m_0, m_{1/2}, A \quad \text{and} \quad \tan\beta \,, \qquad (8)$$

with μ being determined by the electroweak vacuum conditions, up to a sign.

This constrained MSSM (CMSSM) serves
as the basis for the subsequent discussion.

[a] Universality between the squarks and sleptons of different generations is motivated by upper limits on flavour-changing neutral interactions [28], but universality between the soft masses of the L, E^c, Q^c, D^c and U^c is not so well motivated.

It has the merit of being sufficiently specific that the different phenomenological constraints can be combined meaningfully. On the other hand, it is just one of the phenomenological possibilities offered by supersymmetry [29].

4 Constraints on the CMSSM

Important constraints on the CMSSM parameter space are provided by direct searches at LEP and the Tevatron collider [7], as seen in Fig. 4. One of these is the limit $m_{\chi^\pm} \gtrsim 103$ GeV provided by chargino searches at LEP, where the third significant figure depends on other CMSSM parameters. LEP has also provided lower limits on slepton masses, of which the strongest is $m_{\tilde{e}} \gtrsim 99$ GeV, again depending only sightly on the other CMSSM parameters, as long as $m_{\tilde{e}} - m_\chi \gtrsim 10$ GeV. The most important constraints on the u, d, s, c, b squarks and gluinos are provided by the Tevatron collider: for equal masses $m_{\tilde{q}} = m_{\tilde{g}} \gtrsim 300$ GeV. In the case of the $\tilde{t}$, LEP provides the most stringent limit when $m_{\tilde{t}} - m_\chi$ is small, and the Tevatron for larger $m_{\tilde{t}} - m_\chi$. Their effect is almost to exclude the range of parameter space where electroweak baryogenesis is possible [30].

Another important constraint is provided by the LEP limit on the Higgs mass: $m_H > 114.1$ GeV. This holds in the Standard Model, for the lightest Higgs boson h in the general MSSM for $\tan\beta \lesssim 5$, and in the CMSSM for all $\tan\beta$, at least as long as CP is conserved [b]. Since m_h is sensitive to sparticle masses, particularly $m_{\tilde{t}}$, via loop corrections:

$$\delta m_h^2 \propto \frac{m_t^4}{m_W^2} \ln\left(\frac{m_{\tilde{t}}^2}{m_t^2}\right) + \ldots \qquad (9)$$

the Higgs limit also imposes important con-

[b] The lower bound on the lightest MSSM Higgs boson may be relaxed significantly if CP violation feeds into the MSSM Higgs sector [33].



Figure 4. Compilations of phenomenological constraints on the CMSSM for $\tan\beta = 10$ and (a) $\mu > 0$, (b) $\mu < 0$. Representative contours of the selectron, chargino and Higgs masses are indicated, as is the likely physics reach of Run II of the Tevatron Collider in (a). The dark shaded regions are excluded because the LSP is charged, whereas a neutralino LSP has acceptable relic density (10) in the light-shaded regions [31]. The medium-shaded region in (b) is excluded by $b \to s\gamma$ [32].

straints on the CMSSM parameters, principally $m_{1/2}$ as seen in Fig. 4.

Also shown in Fig. 4 is the constraint imposed by measurements of $b \to s\gamma$ [32]. These agree with the Standard Model, and therefore provide bounds on chargino and charged Higgs masses, for example. For moderate $\tan\beta$, the $b \to s\gamma$ constraint is more important for $\mu < 0$, as seen in Fig. 4b, but it is also significant for $\mu > 0$ when $\tan\beta$ is large.

Fig. 4 also displays the regions where the supersymmetric relic density $\rho_\chi = \Omega_\chi \rho_{critical}$ falls within the preferred range

$$0.1 < \Omega_\chi h^2 < 0.3 \qquad (10)$$

The upper limit is rigorous, since astrophysics and cosmology tell us that the total matter density $\Omega_m \lesssim 0.4$, and the Hubble expansion rate $h \sim 1/\sqrt{2}$ to within about 10 % (in units of km/s/Mpc). On the other hand, the lower limit in (10) is optional, since there could be other important contributions to the overall matter density.

As is seen in Fig. 4, there are generic regions of the CMSSM parameter space where the relic density falls within the preferred range (10). What goes into the calculation of the relic density? It is controlled by the annihilation rate [25]:

$$\rho_\chi = m_\chi n_\chi : n_\chi \sim \frac{1}{\sigma_{ann}(\chi\chi \to \ldots)} \qquad (11)$$

and the typical annihilation rate $\sigma_{ann} \sim 1/m_\chi^2$. For this reason, the relic density typically increases with the relic mass, and this combined with the upper bound in (10) then leads to the common expectation that $m_\chi \lesssim 1$ TeV. However, there are various ways in which the generic upper bound on m_χ can be increased along filaments in the $(m_{1/2}, m_0)$ plane. For example, if the next-to-lightest sparticle (NLSP) is not much heavier than χ: $\Delta m/m_\chi \lesssim 0.1$, the relic density may be suppressed by coannihilation: $\sigma(\chi + \mathrm{NLSP} \to \ldots)$ [34]. In this way, the allowed CMSSM region may acquire a 'tail' extending to large m_χ, as in the case where the NLSP is the lighter stau: $\tilde\tau_1$ and $m_{\tilde\tau_1} \sim m_\chi$ as seen in Fig. 5 [35]. Another mechanism for extending the allowed CMSSM region to large m_χ is rapid annihilation via a direct-channel pole when $m_\chi \sim \frac{1}{2} m_{Higgs,Z}$ [36,37]. This may yield a 'funnel' extending to large $m_{1/2}$ and m_0 at large $\tan\beta$, as seen in Fig. 6 [37]. Another allowed region at large $m_{1/2}$ and m_0 is the 'focus-point' region [38], which is adjacent to the boundary of the region where electroweak symmetry breaking is possible, as seen in Fig. 7. However, in this region m_χ is not particularly large.



Figure 5. The large-$m_{1/2}$ 'tail' of the $\chi - \tilde\tau_1$ coannihilation region for $\tan\beta = 10$ and $\mu < 0$ [35].

These filaments extending the preferred CMSSM parameter space are clearly exceptional, in some sense, so it is important to understand the sensitivity of the relic density to input parameters, unknown higher-order effects, etc. One proposal is the relic-density fine-tuning measure [39]

$$\Delta^\Omega \equiv \sqrt{\sum_i \left(\frac{\partial \ln(\Omega_\chi h^2)}{\partial \ln a_i} \right)} \qquad (12)$$



Figure 6. The region where the cosmological relic density is in the preferred range (10) for $\tan\beta = 50$ and $\mu > 0$. Note the rapid-annihilation 'funnel' at intermediate $m_0/m_{1/2}$ [37].



Figure 7. The $m_{1/2}, m_0$ plane for $\tan\beta = 10$ and $\mu > 0$, including the 'focus-point' region [38] at large m_0, close to the boundary of the shaded region where electroweak symmetry breaking occurs, and exhibiting contours of the cosmological sensitivity (12) [39].

where the sum runs over the input parameters, which might include (relatively) poorly-known Standard Model quantities such as m_t and m_b, as well as the CMSSM parameters $m_0, m_{1/2}$, etc. As seen in Fig. 7, the sensitivity Δ^Ω (12) is relatively small in the 'bulk' region at low $m_{1/2}$, m_0, and $\tan\beta$. However, it is somewhat higher in the $\chi - \tilde{\tau}_1$ coannihilation 'tail', and at large $\tan\beta$ in general. The sensitivity measure Δ^Ω (12) is particularly high in the rapid-annihilation 'funnel' and in the 'focus-point' region. This explains why published relic-density calculations may differ in these regions [40], whereas they agree well when Δ^Ω is small: differences may arise because of small differences in the treatments of the inputs.

It is important to note that the relic-density fine-tuning measure (12) is distinct from the traditional measure of the fine-tuning of the electroweak scale [41]:

$$\Delta_i \equiv \frac{\partial \ln m_W}{\partial \ln a_i} \qquad (13)$$

This electroweak fine-tuning is a completely different issue, and values of the Δ_i are not necessarily related to values of Δ^Ω. Electroweak fine-tuning is sometimes used as a criterion for restricting the CMSSM parameters. However, the interpretation of the Δ_i (13) is unclear. How large a value of Δ_i is tolerable? Different physicists may well have different pain thresholds. Moreover, correlations between input parameters may reduce its value in specific models.

5 Muon Anomalous Magnetic Moment

As reported at this meeting [6], the BNL E821 experiment has recently reported a 2.6-σ deviation of $a_\mu \equiv \frac{1}{2}(g_\mu - 2)$ from the Standard Model prediction [42]:

$$a_\mu^{exp} - a_\mu^{th} = (43 \pm 16) \times 10^{-10} \qquad (14)$$

The largest contribution to the error in (14) is the statistical error of the experiment, which will soon be significantly reduced, as many more data have already been recorded. The next-largest error is that due to strong-interaction uncertainties in the Standard Model prediction. Recent estimates converge on an estimate of about 7×10^{-10} for the error in the hadronic vacuum polarization contribution to (14) [43], and the error in the hadron light-by-light scattering contribution is generally thought to be smaller [44]. Therefore, if the central value in (14) does not change substantially with the new data, this would be strong evidence for new physics at the TeV scale.

As many authors have pointed out [45], the discrepancy (14) could well be explained by supersymmetry if $\mu > 0$ and $\tan\beta$ is not too small, as exemplified in Fig. 8. Good consistency with all the experimental and cosmological constraints on the CMSSM is found for $\tan\beta \lesssim 10$ and $m_\chi \simeq 150$ to 350 GeV. Already before the measurement (14), the LHC was thought to have a good chance of discovering supersymmetry [46]. If the result (14) were to be confirmed, this would be almost guaranteed, as we now discuss.



Figure 8. The medium-shaded region is that compatible with the BNL E821 measurement of $g_\mu - 2$ at the 2-σ level [6,42], the light-shaded region has a relic density in the preferred range (10), and the dark-shaded region does not have a neutralino LSP. Good compatibility is found between $g_\mu - 2$ and the other phenomenological constraints for $\tan\beta \sim 5$ or more [45].

6 Prospects for Observing Supersymmetry at Accelerators

As an aid to the assessment of the prospects for detecting sparticles at different accelerators, benchmark sets of supersymmetric parameters have often been found useful [47], since they provide a focus for concentrated discussion. A set of post-LEP benchmark scenarios in the CMSSM has recently been proposed [48], and are illustrated schematically in Fig. 9. They take into account the direct searches for sparticles and Higgs bosons, $b \to s\gamma$ and the preferred cosmological density range (10). About a half of the proposed benchmark points are consistent with $g_\mu - 2$ (14) at the $2-\sigma$ level, but this was not imposed as an absolute requirement.

The proposed points were chosen not to provide an 'unbiased' statistical sampling of the CMSSM parameter space, whatever that means in the absence of a plausible *a priori* measure, but rather are intended to illustrate the different possibilities that are still allowed by the present constraints [48]. Five of the chosen points are in the 'bulk' region at small $m_{1/2}$ and m_0, four are spread along the coannihilation 'tail' at larger $m_{1/2}$ for various values of $\tan\beta$, two are in the 'focus-point' region at large m_0, and two are in rapid-annihilation 'funnels' at large $m_{1/2}$ and m_0. The proposed points range over the allowed values of $\tan\beta$ between 5 and 50. Most of them have $\mu > 0$, as favoured by $g_\mu - 2$, but there are two points with $\mu < 0$.

Various derived quantities in these supersymmetric benchmark scenarios, includ-



Figure 9. Schematic overview of the benchmark points proposed in [48]. They were chosen to be compatible with the indicated experimental constraints, as well as have a relic density in the preferred range (10). The points are intended to illustrate the range of available possibilities.



Figure 10. Estimates of the numbers of different types of CMSSM particles that may be detectable [48] at (a) the LHC, (b) a 1-TeV linear e^+e^- collider [49], and (c,d) a 3(5)-TeV e^+e^- [50] or $\mu^+\mu^-$ collider [51]. Note the complementarity between the sparticles detectable at the LHC and at a 1-TeV linear e^+e^- collider.

ing the relic density, $g_\mu - 2, b \to s\gamma$, electroweak fine-tuning Δ and the relic-density sensitivity Δ^Ω, are given in [48]. These enable the reader to see at a glance which models would be excluded by which refinement of the experimental value of $g_\mu - 2$. Likewise, if you find some amount of fine-tuning uncomfortably large, then you are free to discard the corresponding models.

The LHC collaborations have analyzed their reach for sparticle detection in both generic studies and specific benchmark scenarios proposed previously [46]. Based on these studies, Fig. 10 displays estimates how many different sparticles may be seen at the LHC in each of the newly-proposed benchmark scenarios [48]. The lightest Higgs boson is always found, and squarks and gluinos are usually found, though there are some scenarios where no sparticles are found at the LHC. The LHC often misses heavier weakly-interacting sparticles such as charginos, neutralinos, sleptons and the other Higgs bosons.

The physics capabilities of linear e^+e^- colliders are amply documented in various design studies [49]. Not only is the lightest MSSM Higgs boson observed, but its major decay modes can be measured with high accuracy, as seen in Fig. 11. Moreover, if sparticles are light enough to be produced, their masses and other properties can be measured very precisely, enabling models of supersymmetry breaking to be tested [52].

As seen in Fig. 10, the sparticles visible at an e^+e^- collider largely complement those visible at the LHC [48]. In most of benchmark scenarios proposed, a 1-TeV linear collider would be able to discover and measure precisely several weakly-interacting sparticles that are invisible or difficult to detect at the LHC. However, there are some benchmark scenarios where the linear collider (as well as the LHC) fails to discover supersymmetry. Only a linear collider with a higher centre-of-mass energy appears sure to cover



Figure 11. Analysis of the accuracy with which Higgs decay branching ratios may be measured with a linear e^+e^- collider [53].

all the allowed CMSSM parameter space, as seen in the lower panels of Fig. 12, which illustrate the physics reach of a higher-energy lepton collider, such as CLIC [50] or a multi-TeV muon collider [51].

7 Prospects for Other Experiments

Detection of cold dark matter

Fig. 12 shows rates for the elastic spin-independent scattering of supersymmetric relics [54], including upper limits from the UKDMC, CDMS and Heidelberg experiments [9], as well as the range suggested by the DAMA collaboration [55]. Also shown are the rates calculated in the proposed benchmark scenarios discussed in the previous section, which are considerably below the DAMA range, but may be within reach of future projects. Indirect searches for supersymmetric dark matter via the products of annihilations in the galactic halo or inside the Sun also have prospects in some of the benchmark scenarios [54].

$\mu \to e\gamma$ and related processes



Figure 12. Rates for the elastic spin-independent scattering of supersymmetric relics on (a) protons and (b) neutrons calculated in benchmark scenarios [54], compared with upper limits from the UKDMC, CDMS and Heidelberg experiments [9], as well as the range suggested by the DAMA collaboration [55].



Figure 13. Illustration in one particular lepton-flavour texture model, for $\tan\beta = 30$ and $\mu > 0$, indicating that $\mu \to e\gamma$ decay may occur at a rate close to the present experimental upper limit, in the CMSSM with parameters chosen [57] to match the measured value of $g_\mu - 2$ [6,42].

The BNL E821 report of a possible deviation from the Standard Model suggests that a non-trivial $\mu - \mu - \gamma$ vertex is generated at a scale $\lesssim 1$ TeV. Neutrino oscillations indicate that there are $\Delta L_\mu \neq 0$ processes [8], so it is natural to expect that there might also be a non-trivial $\mu - e - \gamma$ vertex. This is indeed the case in a generic supersymmetric GUT, where neutrino mixing induces slepton mixing [56]. Within this framework, the measurement of $g_\mu - 2$ fixes the sparticle scale, and $\Gamma(\mu \to e\gamma)$ may then be calculated within any given flavour texture. Very approximately, if $g_\mu - 2$ is within one or two σ of the present central value, one may expect $B(\mu \to e\gamma)$ with one or two orders of magnitude of the present experimental upper limit, as illustrated in Fig. 13 [57].

The decay $\mu \to 3e$ and $\mu \to e$ conversion on nuclei are expected to occur with branching ratios within two or three orders of magnitude of $B(\mu \to e\gamma)$, and it is in principle possible to measure CP violation in $\mu \to 3e$ decay. This may provide another interesting interface with neutrino physics and cosmology [58]. The minimal supersymmetric seesaw model has six CP-violating phases: the MNS phase δ, two light-neutrino Majorana phases, and three phases arising from neutrino Dirac Yukawa couplings, which may be responsible for our existence via leptogenesis in the early Universe [8]. The CP-violating neutrino phases induce phases in slepton mass matrices, which may show up in $\mu \to 3e$ decay, $\tau \to 3e/\mu$ decays and leptonic electric dipole moments. In principle, the leptogenesis phases might be obtainable by comparing CP-violating measurements in the charged-lepton and neutrino sectors [58].

Proton decay

This could be within reach, with $\tau(p \to e^+\pi^0)$ via a dimension-six operator possibly $\sim 10^{35} y$ if $m_{GUT} \sim 10^{16}$ GeV as expected in a minimal supersymmetric GUT. Such a model also suggests that $\tau(p \to \bar{\nu}K^+) < 10^{32} y$ via dimension-five operators [59], unless measures are taken to suppress them [60]. This provides motivation for a next-generation megaton experiment that could detect proton decay as well as explore new horizons in neutrino physics [61].

8 Conclusions

As we have seen, future colliders such as the LHC and a TeV-scale linear e^+e^- collider have good prospects of discovering supersymmetry and making detailed measurements. In parallel, B and ν factories have good prospects of making inroads on the flavour and unification problems. Searches for dark matter, stopped-muon experiments and searches for proton decay also have in-

teresting prospects.

Looking further beyond the Standard Model, how can one hope to test a Theory of Everything, including quantum gravity? This should be our long-term ambition, our analogue of the 'faint blue dot' towards which exoplanetary science is directed, and which motivates much of their funding. Testing a quantum theory of gravity will be relatively easy if there are large extra dimensions [10,62]. Much more challenging would be the search for observable effects if the gravitational scale turns out, after all, to be of the same order as the Planck mass $\sim 10^{19}$ GeV. Perhaps the only way to reconcile relativity with quantum mechanics is to modify one or the other, or both [63]? Testing the Theory of Everything may require thinking beyond the standard 'Beyond the Standard Model' box.

References

1. S. Aoki, J. Goodman, C. K. Jung and J. Klein, talks at this conference.
2. M. S. Turner, talk at this conference.
3. R. Barbieri, talk at this conference.
4. M. Neubert, talk at this conference.
5. F. Zwirner, talk at this conference.
6. L. Robertson, talk at this conference.
7. G. Hanson, talk at this conference.
8. H. Murayama, talk at this conference.
9. H. Klapdor-Kleingrothaus, talk at this conference.
10. L. Randall, talk at this conference.
11. D. A. Ross and M. J. Veltman, Nucl. Phys. B **95** (1975) 135.
12. For a classic review,see: E. Farhi and L. Susskind, Phys. Rept. **74** (1981) 277; for a recent review and references, see: K. D. Lane, *Technicolor 2000*, arXiv:hep-ph/0007304.
13. See, for example: G. Altarelli, arXiv:hep-ph/0011078, as updated in G. Altarelli, F. Caravaglios, G. F. Giudice, P. Gambino and G. Ridolfi, JHEP **0106** (2001) 018 [arXiv:hep-ph/0106029].
14. J. R. Ellis, G. L. Fogli and E. Lisi, Phys. Lett. B **343** (1995) 282.
15. J. Drees, talk at this conference. See the LEP Collaborations, ALEPH, DELPHI, L3, OPAL, the LEP Electroweak Working Group and the SLD Heavy Flavour and Electroweak Groups, LEPEWWG/2001-01, available from `http://lepewwg.web.cern.ch/LEPEWWG/Welcome.html`.
16. J. R. Ellis and D. Ross, Phys. Lett. B **506** (2001) 331 [arXiv:hep-ph/0012067].
17. For a review, see: T. Hambye and K. Riesselmann, arXiv:hep-ph/9708416.
18. G. Isidori, G. Ridolfi and A. Strumia, Nucl. Phys. B **609** (2001) 387 [arXiv:hep-ph/0104016].
19. L. Maiani, *Proceedings of the 1979 Gif-sur-Yvette Summer School On Particle Physics*, 1; G. 't Hooft, in *Recent Developments in Gauge Theories, Proceedings of the Nato Advanced Study Institute, Cargese, 1979*, eds. G. 't Hooft *et al.*, (Plenum Press, NY, 1980); E. Witten, Phys. Lett. B **105** (1981) 267.
20. J. Ellis, S. Kelley and D. V. Nanopoulos, Phys. Lett. B **260** (1991) 131; U. Amaldi, W. de Boer and H. Furstenau, Phys. Lett. B **260** (1991) 447; C. Giunti, C. W. Kim and U. W. Lee, Mod. Phys. Lett. A **6** (1991) 1745.
21. K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. B **436** (1998) 55 [arXiv:hep-ph/9803466] and Nucl. Phys. B **537** (1999) 47 [arXiv:hep-ph/9806292].
22. D. M. Ghilencea and G. G. Ross, Nucl. Phys. B **606** (2001) 101 [arXiv:hep-ph/0102306].
23. J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B **257** (1991) 83; M.S. Berger, Phys. Rev. D **41** (1990) 225; Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85** (1991) 1; Phys. Lett.

B **262** (1991) 54; H.E. Haber and R. Hempfling, Phys. Rev. Lett. **66** (1991) 1815; R. Barbieri, M. Frigeni and F. Caravaglios, Phys. Lett. B **258** (1991) 167; P.H. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. B **274** (1992) 191; J.R. Espinosa and M. Quirós, Phys. Lett. B **266** (1991) 389; J.L. Lopez and D.V. Nanopoulos, Phys. Lett. B **266** (1991) 397; M. Carena, K. Sasaki and C.E.M. Wagner, Nucl. Phys. B **381** (1992) 66; P.H. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. B **281** (1992) 100; Nucl. Phys. B **423** (1994) 437; D.M. Pierce, A. Papadopoulos and S.B. Johnson, Phys. Rev. Lett. **68** (1992) 3678; A. Brignole, Phys. Lett. B **281** (1992) 284; M. Drees and M.M. Nojiri, Phys. Rev. D **45** (1992) 2482; V. Barger, M.S. Berger and P. Ohmann, Phys. Rev. D **49** (1994) 4908; G.L. Kane, C. Kolda, L. Roszkowski and J.D. Wells, Phys. Rev. D **49** (1994) 6173; R. Hempfling and A.H. Hoang, Phys. Lett. B **331** (1994) 99; P. Langacker and N. Polonsky, Phys. Rev. D **50** (1994) 2199; H.E. Haber, R. Hempfling and A.H. Hoang, Zeit. für Phys. C **75** (1997) 539; S. Heinemeyer, W. Hollik and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000) [hep-ph/9812320].

24. For a review, see: H. Dreiner, arXiv:hep-ph/9707435.
25. J. Ellis, J.S. Hagelin, D.V. Nanopoulos, K.A. Olive and M. Srednicki, Nucl. Phys. B **238** (1984) 453; see also H. Goldberg, Phys. Rev. Lett. **50** (1983) 1419.
26. S. Dimopoulos and H. Georgi, Nucl. Phys. B **193** (1981) 150; N. Sakai, Z. Phys. C **11** (1981) 153.
27. K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, Prog. Theor. Phys. **68** (1982) 927 [Erratum-ibid. **70** (1982) 330]; L.E. Ibáñez and G.G. Ross, Phys. Lett. B **110** (1982) 215; L.E. Ibáñez, Phys. Lett. B **118** (1982) 73; J. Ellis, D.V. Nanopoulos and K. Tamvakis, Phys. Lett. B **121** (1983) 123; J. Ellis, J. Hagelin, D.V. Nanopoulos and K. Tamvakis, Phys. Lett. B **125** (1983) 275; L. Alvarez-Gaumé, J. Polchinski, and M. Wise, Nucl. Phys. B **221** (1983) 495.
28. J. R. Ellis and D. V. Nanopoulos, Phys. Lett. B **110** (1982) 44; R. Barbieri and R. Gatto, Phys. Lett. B **110** (1982) 211.
29. M. Dine and A. E. Nelson, Phys. Rev. D **48**, 1277 (1993) [hep-ph/9303230]; M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. D **51**, 1362 (1995) [hep-ph/9408384]; M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D **53**, 2658 (1996) [hep-ph/9507378]. D. E. Kaplan, G. D. Kribs and M. Schmaltz, Phys. Rev. D **62**, 035010 (2000) [hep-ph/9911293]; Z. Chacko, M. A. Luty, A. E. Nelson and E. Ponton, JHEP **0001**, 003 (2000) [hep-ph/9911323]. L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999) [hep-th/9810155]; G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998) [hep-ph/9810442].
30. M. Fukugita and T. Yanagida, Phys. Lett. B **174** (1986) 45; for a recent review, see: W. Buchmuller, arXiv:hep-ph/0107153.
31. J. R. Ellis, G. Ganis, D. V. Nanopoulos and K. A. Olive, Phys. Lett. B **502** (2001) 171 [arXiv:hep-ph/0009355].
32. M.S. Alam et al., [CLEO Collaboration], Phys. Rev. Lett. **74** (1995) 2885 as updated in S. Ahmed et al., CLEO CONF 99-10; BELLE Collaboration, BELLE-CONF-0003, contribution to the 30th International conference on High-Energy Physics, Osaka, 2000. See also K. Abe *et al.*, [Belle Collaboration], [arXiv:hep-ex/0107065]; L. Lista [BaBar Collaboration], [arXiv:hep-ex/0110010]; C. Degrassi, P. Gambino and G. F. Giudice,

JHEP **0012** (2000) 009; M. Carena, D. Garcia, U. Nierste and C. E. Wagner, hep-ph/0010003; G. Isidori, talk at this conference.
33. M. Carena, J. R. Ellis, A. Pilaftsis and C. E. Wagner, Nucl. Phys. B **586** (2000) 92 [arXiv:hep-ph/0003180], Phys. Lett. B **495** (2000) 155 [arXiv:hep-ph/0009212]; and references therein.
34. S. Mizuta and M. Yamaguchi, Phys. Lett. B **298** (1993) 120 [arXiv:hep-ph/9208251]; J. Edsjo and P. Gondolo, Phys. Rev. D **56** (1997) 1879 [arXiv:hep-ph/9704361]. C. Boehm, A. Djouadi and M. Drees, Phys. Rev. D **62** (2000) 035012 [arXiv:hep-ph/9911496].
35. J. Ellis, T. Falk and K. A. Olive, Phys. Lett. B **444**, 367 (1998); J. Ellis, T. Falk, K. A. Olive and M. Srednicki, Astropart. Phys. **13** (2000) 181; M. E. Gómez, G. Lazarides and C. Pallis, Phys. Rev. D **61**, 123512 (2000) [hep-ph/9907261] and Phys. Lett. B **487**, 313 (2000) [hep-ph/0004028]; R. Arnowitt, B. Dutta and Y. Santoso, hep-ph/0102181.
36. M. Drees and M. M. Nojiri, Phys. Rev. D **47**, 376 (1993); H. Baer and M. Brhlik, Phys. Rev. D **53** (1996) 597 and Phys. Rev. D **57** (1998) 567; H. Baer, M. Brhlik, M. A. Diaz, J. Ferrandis, P. Mercadante, P. Quintana and X. Tata, Phys. Rev. D **63** (2001) 015007; A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, hep-ph/0009065.
37. J. R. Ellis, T. Falk, G. Ganis, K. A. Olive and M. Srednicki, Phys. Lett. B **510** (2001) 236 [arXiv:hep-ph/0102098].
38. J. L. Feng, K. T. Matchev and T. Moroi, Phys. Rev. Lett. **84**, 2322 (2000) [hep-ph/9908309]; J. L. Feng, K. T. Matchev and T. Moroi, Phys. Rev. D **61**, 075005 (2000) [hep-ph/9909334]; J. L. Feng, K. T. Matchev and F. Wilczek, Phys. Lett. B **482**, 388 (2000) [hep-ph/0004043].
39. J. R. Ellis and K. A. Olive, Phys. Lett. B **514** (2001) 114 [arXiv:hep-ph/0105004].
40. For other recent calculations, see, for example: A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Phys. Lett. B **518** (2001) 94 [arXiv:hep-ph/0107151]; L. Roszkowski, R. Ruiz de Austri and T. Nihei, JHEP **0108**, 024 (2001) [arXiv:hep-ph/0106334].
41. J. Ellis, K. Enqvist, D. V. Nanopoulos and F. Zwirner, Mod. Phys. Lett. A **1**, 57 (1986); R. Barbieri and G. F. Giudice, Nucl. Phys. B **306** (1988) 63.
42. H. N. Brown *et al.* [Muon g-2 Collaboration], Phys. Rev. Lett. **86**, 2227 (2001) [hep-ex/0102017].
43. R. Alemany, M. Davier and A. Hocker, Eur. Phys. J. C **2** (1998) 123 [arXiv:hep-ph/9703220]; M. Davier and A. Hocker, Phys. Lett. B **419** (1998) 419 [arXiv:hep-ph/9711308]; M. Davier and A. Hocker, Phys. Lett. B **435** (1998) 427 [arXiv:hep-ph/9805470]; S. Narison, Phys. Lett. B **513** (2001) 53 [arXiv:hep-ph/0103199]; J. F. De Troconiz and F. J. Yndurain, arXiv:hep-ph/0106025.
44. J. Bijnens, talk at this conference, arXiv:hep-ph/0108111. See also J. Bijnens, E. Pallante and J. Prades, Nucl. Phys. B **474** (1996) 379 [arXiv:hep-ph/9511388]; M. Hayakawa and T. Kinoshita, Phys. Rev. D **57** (1998) 465 [arXiv:hep-ph/9708227].
45. L. L. Everett, G. L. Kane, S. Rigolin and L. Wang, Phys. Rev. Lett. **86**, 3484 (2001) [arXiv:hep-ph/0102145]; J. L. Feng and K. T. Matchev, Phys. Rev. Lett. **86**, 3480 (2001) [arXiv:hep-ph/0102146]; E. A. Baltz and P. Gondolo, Phys. Rev. Lett. **86**, 5004 (2001) [arXiv:hep-ph/0102147]; U. Chattopadhyay and P. Nath, Phys. Rev. Lett. **86**, 5854 (2001) [arXiv:hep-ph/0102157]; S. Komine, T. Moroi and M. Yamaguchi, Phys. Lett. B **506**, 93 (2001) [arXiv:hep-ph/0102204]; S. P. Martin and J. D. Wells, Phys. Rev. D **64**,

035003 (2001) [arXiv:hep-ph/0103067]; H. Baer, C. Balazs, J. Ferrandis and X. Tata, Phys. Rev. D **64**, 035004 (2001) [arXiv:hep-ph/0103280]; J. Ellis, D. V. Nanopoulos and K. A. Olive, Phys. Lett. B **508** (2001) 65 [arXiv:hep-ph/0102331]; R. Arnowitt, B. Dutta, B. Hu and Y. Santoso, Phys. Lett. B **505** (2001) 177 [arXiv:hep-ph/0102344].

46. ATLAS Collaboration, *ATLAS detector and physics performance Technical Design Report*, CERN/LHCC 99-14/15 (1999); S. Abdullin *et al.* [CMS Collaboration], hep-ph/9806366; S. Abdullin and F. Charles, Nucl. Phys. B **547** (1999) 60; CMS Collaboration, Technical Proposal, CERN/LHCC 94-38 (1994).
47. See, for example: I. Hinchliffe, F. E. Paige, M. D. Shapiro, J. Soderqvist and W. Yao, Phys. Rev. D **55** (1997) 5520; TESLA Technical Design Report, DESY-01-011, Part III, *Physics at an e^+e^- Linear Collider* (March 2001).
48. M. Battaglia *et al.*, arXiv:hep-ph/0106204.
49. S. Matsumoto *et al.* [JLC Group], *JLC-1*, KEK Report 92-16 (1992); J. Bagger *et al.* [American Linear Collider Working Group], *The Case for a 500-GeV e^+e^- Linear Collider*, SLAC-PUB-8495, BNL-67545, FERMILAB-PUB-00-152, LBNL-46299, UCRL-ID-139524, LBL-46299, Jul 2000, hep-ex/0007022; T. Abe *et al.* [American Linear Collider Working Group Collaboration], *Linear Collider Physics Resource Book for Snowmass 2001*, SLAC-570, hep-ex/0106055, hep-ex/0106056, hep-ex/0106057 and hep-ex/0106058; TESLA Technical Design Report, DESY-01-011, Part III, *Physics at an e^+e^- Linear Collider* (March 2001).
50. R. W. Assmann *et al.* [CLIC Study Team], *A 3-TeV e^+e^- Linear Collider Based on CLIC Technology*, ed. G. Guignard, CERN 2000-08; CLIC Physics Study Group, `http://clicphysics.web.cern.ch/CLICphysics/`.
51. Neutrino Factory and Muon Collider Collaboration, `http://www.cap.bnl.gov/mumu/mu_home_page.html`; European Muon Working Groups, `http://muonstoragerings.cern.ch/Welcome.html`.
52. G. A. Blair, W. Porod and P. M. Zerwas, Phys. Rev. **D63** (2001) 017703 [hep-ph/0007107].
53. M. Battaglia and K. Desch, arXiv:hep-ph/0101165.
54. J. Ellis, J. L. Feng, A. Ferstl, K. T. Matchev and K. A. Olive, arXiv:astro-ph/0110225.
55. DAMA Collaboration, R. Bernabei *et al.*, Phys. Lett. B **436**, 379 (1998).
56. J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D **53** (1996) 2442; J. Hisano, D. Nomura and T. Yanagida, Phys. Lett. B **437** (1998) 351; J. Hisano and D. Nomura, Phys. Rev. D **59** (1999) 116005; W. Buchmüller, D. Delepine and F. Vissani, Phys. Lett. B **459** (1999) 171; M. E. Gomez, G. K. Leontaris, S. Lola and J. D. Vergados, Phys. Rev. D **59** (1999) 116009; J. R. Ellis, M. E. Gomez, G. K. Leontaris, S. Lola and D. V. Nanopoulos, Eur. Phys. J. C **14** (2000) 319; W. Buchmüller, D. Delepine and L. T. Handoko, Nucl. Phys. B **576** (2000) 445; J. L. Feng, Y. Nir and Y. Shadmi, Phys. Rev. D **61** (2000) 113005; J. Sato and K. Tobe, Phys. Rev. D **63** (2001) 116010; J. Hisano and K. Tobe, Phys. Lett. B **510** (2001) 197; S. Baek, T. Goto, Y. Okada and K. Okumura, hep-ph/0104146; S. Lavignac, I. Masina and C.A. Savoy, hep-ph/0106245.
57. D. F. Carvalho, J. R. Ellis, M. E. Gomez and S. Lola, Phys. Lett. B **515** (2001)

323 [arXiv:hep-ph/0103256].
58. J. R. Ellis, J. Hisano, S. Lola and M. Raidal, arXiv:hep-ph/0109125; and references therein.
59. H. Murayama and A. Pierce, arXiv:hep-ph/0108104.
60. J. R. Ellis, J. S. Hagelin, S. Kelley and D. V. Nanopoulos, Nucl. Phys. B **311** (1988) 1.
61. C. K. Jung, arXiv:hep-ex/0005046; Y. Suzuki *et al.* [TITAND Working Group Collaboration], arXiv:hep-ex/0110005.
62. I. Antoniadis and K. Benakli, Int. J. Mod. Phys. A **15** (2000) 4237 [arXiv:hep-ph/0007226].
63. J. R. Ellis, N. E. Mavromatos and D. V. Nanopoulos, arXiv:gr-qc/9909085.